

---

# RELATIONSHIP BETWEEN BODY MASS AND BODY LENGTH IN CAPYBARAS (*HYDROCHOERUS HYDROCHAERIS*)

*Katia Maria Paschoaletto Micchi de Barros Ferraz, Kelly Bonach, Luciano Martins Verdade*

Biota Neotropica v5(n1) – <http://www.biotaneotropica.org.br/v5n1/pt/abstract?short-communication+BN03405012005>

*Recebido em 10/12/2004*

*Revisado em 10/20/2005*

*Publicado 14/03/2005*

Laboratório de Ecologia Animal, ESALQ, Universidade de São Paulo  
Cx. P. 09, Piracicaba, SP, 13418-900 BRASIL  
Email: [kferraz@esalq.usp.br](mailto:kferraz@esalq.usp.br)

## Abstract

As a part of a management program we captured 39 capybaras (*Hydrochoerus hydrochaeris*) in an agroecosystem at the east central region of the State of São Paulo, Brazil from March 2001 to May 2002. Average adult male body mass was  $54.1 \pm 8.05$  kg, and average adult female body mass was  $62.0 \pm 12.03$  kg. Average juvenile male body mass was  $23.0 \pm 8.28$  kg, and average juvenile female body mass was  $26.7 \pm 5.86$  kg. Males and females presented a significant variation in the allometric relation between body mass and body length. Body mass and body length had a high correlation for both adult males and adult females. These results from capybaras in agroecosystem might be relevant for further management programs.

**Key words:** *Capybaras, body mass, body length, agroecosystem*

## Resumo

Trinta e nove capivaras (*Hydrochoerus hydrochaeris*) foram capturadas como parte de um programa de manejo em um agroecossistema na região centro-oeste do estado de São Paulo, de março de 2001 a maio de 2002. A massa corpórea média de machos adultos foi de  $54.1 \pm 8.05$  kg, enquanto que de fêmeas adultas foi de  $62.0 \pm 12.03$  kg. A massa corpórea média de machos jovens foi de  $23.0 \pm 8.28$  kg, enquanto que de fêmeas jovens foi de  $26.7 \pm 5.86$  kg. Machos e fêmeas apresentaram variação significativa na relação alométrica entre massa e comprimento corpóreos. Massa e comprimento corpóreos apresentaram alta correlação entre machos e fêmeas adultas. Estes resultados de capivaras em agroecossistemas podem ser importantes para futuros programas de manejo.

**Palavras-chave:** *capivara, massa corpórea, comprimento corpóreo, agroecossistemas*

## Introduction

Capybaras, the largest living rodent in the world, have been considered as a potential plague in the east central region of the State of São Paulo, especially in the Piracicaba river basin, a well developed and anthropogenic region in the State. Apparently, the process of habitat deforestation has been favoring capybaras population increase in this region, mainly due to the great food availability by agriculture such as corn, sugar cane, and rice (Ferraz et al., 2003). Besides the natural vegetation (grass and aquatic plants - Ojasti, 1973), capybaras consume mainly C4 domestic plants what has been probably contributing for the species adaptation to anthropogenic habitats. In this region, capybara habitats can be described as agricultural fields in flat areas near watercourses (Ferraz 2004). Capybara has also been reported to cause damage in agricultural fields at the University Campus (Ferraz et al., 2003) and its relationship with the occurrence of rickettsiosis has also been studied (Labruna et al., 2004). As part of a management program, capybaras have been monitored in this study sites weekly by direct counts since July 1998 (Verdade & Ferraz 2005). In this study we present some body mass and body length from 39 capybaras captured in these anthropogenic habitats.

## Material and Methods

The site where capybaras were captured could be described as an agroecosystem located at Campus "Luiz de Queiroz", from the University of São Paulo, in Piracicaba, Brazil (22°41'51.2" S and 47°33'40.0" W, in average 546 m above sea level) (Figure 1). The ecosystem was composed by a mixed field of agriculture (mainly corn, approximately 90 ha), a fragment of second-growth semi-deciduous native forest (7 ha), a stand of mixed implanted forest (*Pinus* spp and *Eucalyptus* spp, 2.7 ha), a small field of exotic pasture (mostly *Brachiaria decumbens*, approximately 1 ha), and an artificial reservoir (1.7 ha).




Figure 1 - Location of the study area, Piracicaba, São Paulo, Brazil.

As part of a management program we captured 39 capybaras (*Hydrochoerus hydrochaeris*) in this agroecosystem from March 2001 to May 2002 (n=10 in March 2001; n=28 in October 2001; and n=1 in May 2002) in two small-fenced area. Based on the species monitoring data, we believe that all captured individuals belonged to the same group.

## Results and Discussion

Fifteen adults (nine males and six females), 19 juveniles (five males and fourteen females), and five young (two males and three females) were captured in the study area. Average adult male body mass was  $54.1 \pm 8.05$  kg (44.0-68.0) and average adult female body mass was  $62.0 \pm 12.03$  kg (48.0-81.0). Average juvenile male body mass was  $23.0 \pm 8.28$  kg (15.0-37.0) and average juvenile female body mass was  $26.7 \pm 5.86$  kg (15.0-34.0) (Table 1). The average body mass of adults and juveniles males and females from the present study was bigger than those from other studies (Ojasti 1973, Cordero & Ojasti 1981, Schaller 1983, Herrera 1986, Mones & Ojasti 1986, Herrera 1992), except from Alho et al. (1989) (Table 2). The present results suggest that in agroecosystems capybaras can reach bigger body mass than in natural habitats possibly because of food availability (Ferraz et al. 2003).

Body mass can be an important attribute to define social position in the dominance hierarchy of capybara grouping. According to Moreira et al. (1997), body mass was significantly correlated to the volume of the snout and anal gland, presence to the coated hairs in the anal glands and testes mass, suggesting that dominance in capybaras is more associated with androgen-dependent scent glands than with sperm production. Mones & Ojasti (1986) suggest that capybaras evince a latitudinal cline with body size and mass increasing as latitude increases.

The heaviest individual found at the present study was an 81.0 kg non-pregnant female, close to the record described by Mones (1973) of a 91 kg female also found in São Paulo, Brazil, but above the heaviest female (65.5 kg) recorded by Ojasti (1973) in Venezuela. At this study most females were non-pregnant contrary to the pattern found by Ojasti (1973). These different patterns could be possibly explained by the seasons when captures were made in both studies. In São Paulo, there seems to be a peak at the reproductive activity in November (Verdade & Ferraz, 2005). Cordero & Ojasti (1981) assumed that a population of capybaras in a forested habitat presented older and therefore heavier individuals than another population from Savannah habitat.

In this study, males and females presented similar body mass ( $t = 0.90$ , d.f. = 32,  $P = 0.37$  for the whole sample,  $t = 1.54$ , d.f. = 13,  $P = 0.14$  for the adults, and  $t = 1.09$ , d.f. = 17,  $P = 0.29$  for the juveniles).

TABLE 1. *Capibaras' body mass (mean + standard error of mean) of males and females, adults and juveniles.*

	<b>Body mass (kg)</b>	<b>Min.</b>	<b>Max.</b>	<b>n</b>
Males (adults and juveniles)	42.96 ± 17.30	15.0	68.0	14
Females (adults and juveniles)	41.73 ± 17.36	15.0	81.0	20
Adults (males and females)	57.23 ± 10.24	44.0	81.0	15
Juveniles (males and females)	25.74 ± 6.54	15.0	37.0	19
Adult males	54.06 ± 8.05	44.0	68.0	9
Adult females	62.00 ± 12.03	48.0	81.0	6
Juvenile males	23.00 ± 8.28	15.0	37.0	5
Juvenile females	26.71 ± 5.86	15.0	34.0	14

TABLE 2. *Body mass (kg) of capybaras.*

Localities	Average body mass (kg)				References
	Adult male	Adult female	Juvenile male	Juvenile female	
São Paulo, Brazil	54.1	62.0	23.0	26.7	Present study
Pantanal, Brazil	41.8	36.8	-	-	Schaller (1983)
	57.57	51.19	14.08	15.55	Alho et al. (1989)
Llanos, Venezuela	48.63	49.80	-	-	Ojasti (1973)
	38.4 - 42.4	37.9 - 39.9	-	-	Herrera (1986)
	43.1	42.2	13.9	14.3	Herrera (1992)

Likewise, males and females presented similar body length (from the top of the nose to the end of the tail) ( $t = 0.89$ , d.f. = 23,  $P = 0.38$  for the whole sample,  $t = -0.07$ , d.f. = 17,  $P = 0.94$  for the adults, and  $t = -1.46$ , d.f. = 12,  $P = 0.17$  for the juveniles). However, males and females presented a significant variation (ANCOVA:  $F = 5.05$ , d.f. = 24,  $P = 0.035$ ) in the allometric relation between body mass and body length. Therefore, we can infer that males and females present similar body size, but different body shape, what means that capybaras present some sexual dimorphism at least in adults. Herrera (1986) noticed a slight sexual dimorphism in adults in the relationship between their body mass and length of hind foot.

Body mass and body length had a high correlation for both adult males and adult females ( $R^2 = 0.98$  and  $R^2 = 0.94$ , respectively). The relationship between body mass (kg) and body length (cm) for adult can be described by the following regression models:  $BM_m = -73.8256 + 1.03223 L_m$  ( $R^2 = 97.9\%$ ), and  $BM_f = -81.1077 + 1.07097 L_f$  ( $R^2 = 94.0\%$ ), where  $BM_m$  = male body mass (kg),  $BM_f$  = female body mass (kg),  $L_m$  = male length (cm), and  $L_f$  = female length (cm). Although the results presented here are related to a small sample size, they are the first records from capybaras of an anthropogenic habitat. These results might be relevant for further management programs.

### Acknowledgments

The present study has been partially supported by Fundação de Amparo à Pesquisa do Estado de São Paulo – FAPESP (Procs. Nos. 00/00180-9, 00/00602-0, and 00/00582-0) and Prefeitura do Campus “Luiz de Queiroz” / USP. Gustavo R. M. Pinto, Marli Penteadó and Paulo Bezerra participated in the fieldwork.

### Literature Cited

- ALHO, C.J.R., CAMPOS, Z.M. & GONÇALVES, H.C. 1989. Ecology, social behavior and management of the capybara in the Pantanal of Brazil. In: *Advances in Neotropical Mammalogy* (Redford, K.H. & Eisenberg, J.F., eds.). Sandhill Crane, Gainesville, FL. p.163-194.
- CORDERO R., G.A., OJASTI, J. 1981. Comparison of capybara populations of open and forested habitats. *J. Wildlife Manage* 45(1): 267-271.
- FERRAZ, K.M.P.B. 2004. Distribuição espacial da capivara (*Hydrochoerus hydrochaeris*) em função da paisagem na bacia do rio Piracicaba, SP. Tese (doutorado). Escola Superior d'Agricultura “Luiz de Queiroz”, Universidade de São Paulo. 115p.
- FERRAZ, K.M.P.M.B., LECHEVALIER, M., COUTO, H.T.Z. & VERDADE, L.M. 2003. Damage caused by capybaras in a corn field, São Paulo, Brasil. *Scientia Agricola* 60(1): 191-194.
- HERRERA, E.A. 1986. The behavioural ecology of capybara, *Hydrochoerus hydrochaeris*. Ph.D. Thesis, University of Oxford, Oxford, U.K.
- HERRERA, E.A. 1992. Growth and dispersal of capybaras (*Hydrochoerus hydrochaeris*) in the Llanos of Venezuela. *J. Zool. London* 228: 307-316.
- LABRUNA, M.B., WHITWORTH, T., HORTA, M.C., BOUYER, D.H., MCBRIDE, J.W., PITER, A., POPOV, V., GENNARI, S.M. & WALKER, D.H. 2004. *Rickettsia* species infecting *Amblyomma cooperi* ticks from an area in the state of São Paulo, Brazil, where Brazilian Spotted Fever is endemic. *J Clin Microbiology* 42(1): 90-98.
- MONES, A. 1973. Estudios sobre la familia Hydrochoeridae (Rodentia), I. Introducción e historia taxonômica. *Rev. Bras. Biol* 33: 277-283.
- MONES, A., OJASTI, J. 1986. *Hydrochoerus hydrochaeris*. *Mamm Species* 264: 1-7.
- MOREIRA, J.R., MACDONALD, D.W. & CLARKE, J.R. 1997. Correlates of testis mass in capybaras (*Hydrochoerus hydrochaeris*): dominance assurance or sperm production. *J. Zool. London* 241: 457-463.
- OJASTI, J. 1973. *Estudio Biológico del Chigüire o Capibara*. Fondo Nacional de Investigaciones Agropecuarias, Caracas, Venezuela.
- SCHALLER, G.S. 1983. Mammals and their biomass on a Brazilian ranch. *Arquiv Zool* 31(1):1-36.
- VERDADE, L.M. & FERRAZ, K.M.P.M.B. In press. Capybaras on an anthropogenic habitat in Southeastern Brazil. *Brazilian Journal of Biology* 66(2) (in press)

Title: Relationship between body mass and body length in Capybaras (*Hydrochoerus hydrochaeris*).

Authors: Katia Maria Paschoaletto Micchi De Barros Ferraz, Kelly Bonach, Luciano Martins Verdade

Biota Neotropica, Vol. 5 (number 1): 2005  
<http://www.biotaneotropica.org.br/v5n1/pt/abstract?short-communication+BN03405012005>

Recebido em 10/12/2004  
 Revisado em 10/20/2005  
 Publicado 14/03/2005

ISSN 1676-0611